### Defining the Magnitude: Patterns, Regularities and Direct TOA-Surface Flux Relationships in the 15-Year Long CERES Satellite Data — Observations, Model and Theory

Miklos Zagoni<sup>1</sup>

<sup>1</sup>Affiliation not available

November 21, 2022

#### Abstract

Over the past fifteen years, the NASA Clouds and the Earth's Radiant Energy System (CERES) satellite mission has provided the scientific community with the most reliable Earth radiation budget data. This presentation offers quantitative assessment of the published CERES Energy Balanced and Filled (EBAF) Edition 2.8 and Edition 4.0 data products, and reveals several internal patterns, ratios and regularities within the annual global mean flux components of the all-sky and clear-sky surface and atmospheric energy budgets. The found patterns, among others, include: (i) direct relationships between the top-of-atmosphere (TOA) radiative and surface radiative and non-radiative fluxes (contradicting the expectation that TOA and surface fluxes are physically decoupled); (ii) integer ratios and relationships between the absorbed and emitted surface and atmospheric energy flow elements; and (iii) definite connections among the clear-sky and the all-sky shortwave, longwave and non-radiative (turbulent) flux elements and the corresponding greenhouse effect. Comparison between the EBAF Ed2.8 and Ed4.0 SFC and TOA data products and trend analyses of the normalized clear-sky and all-sky greenhouse factors are presented. Longwave cloud radiative effect (LW CRE) proved to be playing a principal role in organizing the found numerical patterns in the surface and atmospheric energy flow components. All of the revealed structures are quantitatively valid within the one-sigma range of uncertainty of the involved individual flux elements. This presentation offers a conceptual framework to interpret the found relationships and shows how the observed CERES fluxes can be deduced from this proposed physical model. An important conclusion drawn from our analysis is that the internal atmospheric and surface energy flow system forms a definite structure and seems to be more constrained to the incoming solar energy than previously thought.

All-sky CERES	EBAF Ed 4.0	EBAF Ed 2.8	Edition M7	N	Ed 4.0 – Ed M7	Ed 2.8 – Ed M7	Ed 4.0 -
TOA LW	240.1	239.6	240.1	9	0.0	-0.5	0.5
SFC SW net	163.7	162.3	160.1	6	3.6	2.2	1.4
SFC LW down	345.0	345.2	346.8	13	-1.8	-1.6	-0.2
SFC SW+LW	508.7	507.5	506.9	19	1.8	0.6	1.2
SFC LW up	398.3	398.3	400.2	15	-1.9	-1.9	0.0
SFC SW+LW net	110.3	109.2	106.7	4	3.6	2.5	1.1
20LR + LWCRE	511.1	508.1	506.9	19	4.2	1.2	3.0
G	158.2	158.7	160.1	6	-1.9	-1.4	-0.5
Clear-sky CERES							
TOA LW	268.1	265.4	266.8	10	1.3	-1.4	2.7
SFC SW net	213.9	214.3	213.4	8	0.5	0.9	0.4
SFC LW down	314.1	316.3	320.2	12	-6.1	4.1	-2.2
SFC SW+LW	528.0	530.6	533.6	20	-5.6	-3.0	-2.6
SFC LW up	397.6	398.4	400.2	15	-2.6	-1.8	-0.8
SFC SW+LW net	130.4	132.2	133.4	5	-3.0	-1.2	-1.8
20LR	536.2	530.8	533.6	20	2.6	-2.8	5.4
G	129.5	133.0	133.4	5	-3.9	-0.4	-3.5
LW CRE							
TOA	28.0	25.8	26.68	1	1.3	-0.9	3.2
SFC	30.9	28.9	26.68	1	4.2	2.2	2.0

#### Defining the Magnitude: Patterns, Regularities and Direct TOA-Surface Flux Relationships A53G-2367 in the 15-Year Long CERES Satellite Data — Observations, Model and Theory

Miklós Zágoni, Eötvös Loránd University; email:miklos.zagoni@t-online.hu

#### Observations

CERES EBAF TOA and SFC Ed2.8 and Ed4.0 data Global Means (Mar2000-Feb2016) (Rose et al. 2017)

				, (		
All S	iky	Ed4	Ed2.8	Clear Sky	Ed4	Ed2.8
TOA	SW Insolation	340.04	339.87	TOA SW Insolation	340.04	339.87
ΤΟΑ	SW Up	99.23	99.62	TOA SW Up	53.41	52.50
ΤΟΑ	LW Up	240.14	239.60	TOA LW Up	268.13	265.59
SFC	SW Down	187.04	186.47	SFC SW Down	243.72	244.06
SFC	SW Up	23.37	24.13	SFC SW Up	29.81	29.74
SFC	LW Down	344.97	345.15	SFC LW Down	314.07	316.27
SFC	LW Up	398.34	398.27	SFC LW Up	397.59	398.40

#### There are patterns in these fluxes

Pattern 1: The SFC absorbed energy (SW down - SW up + LW down) in the clear-sky equals to 2OLR, with a difference of 0.6 Wm<sup>-2</sup> in Ed2.8 and 8.3 Wm<sup>-2</sup> in Ed4.0. Pattern 2: The net (non-radiative) energy at the surface in clear-sky = SFC Net = SW down - SW up + LW down - LW up equals to the G greenhouse effect = SFC LW up - TOA LW up with Diff = -0.6 Wm-2 (Ed2.8) and 0.9 Wm-2 (Ed4.0) Pattern 3: Ed 2.8 Clear-sky integer ratios: G = SFC LW up - TOA LW up =132.8 (1); TOA LW up = 265.6 (2), SFC LW up = 398.4 (3); E(SFC) = 530.6 (4), Pattern 4: Pattern 1 is valid for the all-sky by adding surface LW cloud radiative effect SW net + LW down = 20LR + SFC LWCRE. Diff = 0.7 (Ed2.8), 0.9 (Ed4) Wm<sup>-2</sup> Pattern 5: All-sky integer ratios: with UNIT = OLR(all-sky)/9 = 240.1/9 = 26.68 W/m<sup>2</sup> (Ed4)

#### Clear-sky, Ed2.8, time period 2001-2015, climate year:

	SW down	SW up	SW abs	LW abs	E(SFC)	OLR	20LR	Diff
CLIM 1	255.35	31.34	224.01	306.27	530.28	262.43	524.86	5.42
CLIM 2	251.86	29.86	222	307.98	529.98	262.78	525.56	4.42
CLIM 3	246.89	30.09	216.8	311.09	527.89	263.42	526.84	1.05
CLIM 4	242.26	31.63	210.63	315.14	525.77	265.01	530.02	-4.2
CLIM 5	237.33	32.21	205.12	320.27	525.39	267.18	534.36	-8.9
CLIM 6	233.41	28.67	204.74	325.70	530.44	269.05	538.1	-7.6
CLIM 7	231.63	25.72	205.91	328.35	534.26	269.75	539.5	-5.2
CLIM 8	233.65	24.42	209.23	327.05	536.28	269.12	538.24	-1.9
CLIM 9	239.24	25.80	213.44	321.82	535.26	267.58	535.16	0.1
CLIM 10	247.06	29.92	217.14	315.28	532.42	265.25	530.5	1.92
CLIM 11	253.97	35.15	220.24	309.57	529.81	20.5.20	526.52	3.25
CLIM 12	256.35	33.52	222.83	306.74	529.57	262.38	524.76	4.81
Average	244.08	29.74	214.34	316.27	530.61	265.60	531.20	-0.5

214.34 +316.27 = 530.61 = 2 × 265.60 - 0.59 Pattern 1

Wild et al. (2016

79.8

3

Clear-sky emissio

266.4±3.

266.0 = 10

399.0 = 15

Diff = 0.4 W/

Cle	ear-sk	y, Ed2	.8, clii	mate ye	ear, 200	01-20	15
	ULW	OLR	G	Net SFC	ULW+G	20LR	Diff
CLM 1	399.3	262.43	125.87	141.08	514.17	524.86	10.69
CLIM 2	389.89	262.78	127.11	140.09	517	525.56	8.56
CLIM 3	393.31	263.42	129.89	134.57	523.2	526.84	3.64
CLIM 4	398.51	265.01	133.5	127.26	532.01	530.02	-1.99
CLIM 5	403.29	267.18	136.11	122.09	539.4	534.36	-5.04
CLIM 6	407.64	269.05	138.59	122.8	546.23	538.1	-8.13
CLIM 7	409.1	269.75	139.35	125.17	548.45	539.5	-8.95
CLIM 8	407.83	269.12	138.71	128.46	546.54	538.24	-8.30
CLIM 9	403.85	267.58	136.27	131.41	540.12	535.16	-4.96
CLIM 10	397.76	265.25	132.51	134.65	530.27	530.5	0.23
CLIM 11	392.27	263.26	129.01	137.54	521.28	526.52	5.24
CLIM 12	389.00	262.38	126.62	140.57	515.62	524.76	9.14
average	398.39	265.60	132.80	132.22	531.19	531.20	0.01

ULW – G = OLR (def.); Data: G = Net SFC (= SH+LH) ULW + G = 20LR, Diff = 0.01 (!!!) W/m<sup>2</sup> =>

G = 0	Patte	ern 2					
Pattern	4 SW abs	LW abs	E(SFC)	OLR(all)	SFC LWCRE	20LR+LWCRE	Diff
CLIM1	166.42	335.38	501.8	236.84	29.49	503.17	-1.37
CLIMZ	167.56	337.14	504.7	237.41	29.51	504.33	U.37
CLIM3	166.66	339.93	506.59	237.72	28.89	504.33	2.26
CLIM4	162.97	344.67	507.64	238.42	29.58	506.42	1.22
CLIM5	157.91	349.22	507.13	240.24	28.64	509.12	-1.99
CLIM6	155.39	353.4	508.79	242.38	28.21	512.97	-4.18
CLIM7	156.73	355.86	512.59	243.71	28.16	515.58	-2.99
CLIM8	160.33	355.23	515.56	243.68	28.2	515.56	0
CLIM9	163.59	350.78	514.37	242.13	28.43	512.69	1.68
CLIM10	164.08	345.39	509.47	239.66	28.98	508.3	1.17
CLIM11	163.66	339.84	503.5	237.19	29.82	504.2	-0.7
CLIM12	164.19	336.4	500.59	236.31	29.75	502.37	-1.78
Average	162.46	345.27	507.73	239.64	28.97	508.25	-0.53
SFC	(SW in	+ LW	in) = 2	OLR(a	II) + SFC LW	CRE (-0.53	W/m²)

#### Model

The observed CERES EBAF fluxes can be modeled as integer multiples of a unit flux. The model data set is Edition MZ.

All-sky	Ed4.0	Ed2.8	EdMZ	N	Ed4.0 – EdMZ	Ed2.8 – EdMZ
TOA SW In	340.04	339.87	340.04		0.0	-0.27
TOA LW Up	240.14	239.60	240.14	9	0.0	-0.54
SFC SW In	163.67	162.34	160.09	6	3.58	2.25
SFC LW Down	344.97	345.15	346.87	13	-1.90	-1.72
SFC (SW in + LW in)	508.64	507.49	506.96	19	1.68	0.53
SFC LW Up	398.34	398.27	400.23	15	-1.89	-1.96
SFC Net	110.30	109.22	106.73	4	3.57	2.49
G	158.20	158.67	160.09	6	-1.89	-1.42
2TOA LW Up+LWCRE	511.18	508.08	506.96	19	4.22	1.12

Clear-sky	Ed4.0	Ed2.8	EdMZ	Ν	Ed4.0 -	Ed2.8 -
					EdMZ	EdMZ
TOA SW in	340.04	339.87	340.04		0.0	-0.17
TOA LW up	268.13	265.59	266.82	10	1.31	-1.23
SFC SW in	213.91	214.32	213.47	8	0.44	0.85
SFC LW in	314.07	316.27	320.18	12	-6.11	-3.91
SFC SW + LW in	527.98	530.59	533.65	20	-5.67	-3.06
SFC LW up	397.59	398.40	400.23	15	-2.64	-1.83
SFC Net	130.39	132.19	133.42	5	-3.03	-1.23
G	129.46	132.81	133.42	5	-3.96	-0.61



#### Theory

The simplest greenhouse model: A planet surrounded by an SW-transparent, LW-opaque, non-turbulent "glass shell" atmosphere. Here the total surface absorption is exactly twice the outgoing longwave radiation, because of the geometry.



From here, all the patterns, including the integer ratios, can be deduced by using simple arithmetic relationships. Clear-sky integer ratios in Costa-Shine (2012) LBL-computation:



#### Conclusions

There are robust patterns in the CERES global means. What are they: coincidences? Or conspiracy 🕲 ? No; they have sound theoretical basis: integer data set is presented to reproduce the patterns, and a simple greenhouse model (IR-opaque limit, glass shell geometry) offers the physical background for the basic ratios. All fluxes are within 1 $\sigma$  range. Compared to the model data, the difference to Ed4.0 clear-sky SFC SW absorption is only 0.44 Wm<sup>-2</sup> and to all-sky SFC LW down is -1.9 Wm<sup>-2</sup>; the clear-sky DLR in Ed4.0 is too low by 6.1 Wm<sup>-2</sup> and the clear-sky OLR is too high by 1.3 Wm<sup>-2</sup>.

http://www.globalenergybudget.com

#### Poster:

AGU 2017 Fall Meeting: A53G-2367 Defining the Magnitude: Patterns, Regularities and Direct TOA-Surface Flux Relationships in the 15-Year Long CERES Satellite Data — Observations, Model and Theory

#### ePoster:

https://agu.confex.com/agu/fm17/meetingapp.cgi/Paper/205884 website: www.globalenergybudget.com email: miklos.zagoni@t-online.hu

#### Handout:

28<sup>th</sup> CERES Science Team Meeting Presentation 2017 September 27, NASA Goddard Space Flight Center Greenbelt, MD.

Miklos Zagoni

# Patterns in the CERES Global Mean Data

Land + Sea solar reflected TOA 100 (340, 341) 2 100 (95 100) 100 (95 100)
3 Miklos Zagoni 1   Internet Internet Internet
Fall 2017 CERES Science Team Meeting, September 27, NASA GSFC, Greenbelt, MD.

"To search for something – though it be mushrooms – or some pattern – is impossible, unless you look and try." Dmitri Mendeleev

# CERES\_EBAF-Surface\_Ed4.0 Data Quality Summary (May 26, 2017)

Table 5-1. Global mean surface fluxes in W m<sup>-2</sup> computed from EBAF Ed4.0 and EBAF Ed2.8 for March 2000-February 2016.

All-sky	Ed4	Ed2.8	Ed4 – Ed2.8
TOA SW insolation	340.0	339.9	0.17
SW down	187.0	186.5	0.57
SW up	23.4	24.1	-0.76
SW net <sup>1</sup>	163.7	162.3	1.33
LW down	345.0	345.2	-0.18
LW up	398.3	398.3	0.07
LW net <sup>1</sup>	-53.4	-53.1	-0.25
SW+LW net	110.3	109.2	1.08
Clear-sky			
TOA SW insolation	340.0	339.9	0.17
SW down	243.7	244.1	-0.33
SW up	29.8	29.7	0.07
SW net <sup>1</sup>	213.9	214.3	0.41
LW down	314.1	316.3	-2.20
LW up	397.6	398.4	-0.81
LW net <sup>1</sup>	-83.5	-82.1	1.39
SW+LW net <sup>1</sup>	130.4	132.2	-1.80

#### Fred Rose et al Global Means(Mar2000-Feb2016)

	and the second		
All Sky	Ed4	Ed2.8	Ed4 –Ed2.8
TOA SW Insolation	340.04	339.87	0.17
TOA SW Up	99.23	99.62	-0.39
TOA LW Up	240.14	239.60	0.54
SFC SW Down	187.04	186.47	0.57
SFC SW Up	23.37	24.13	-0.76 (3.1%)
SFC LW Down	344.97	345.15	-0.18
SFC LW Up	398.34	398.27	0.07
Clear Sky	Ed4	Ed2.8	Ed4 –Ed2.8
Clear Sky TOA SW Insolation	Ed4 340.04	Ed2.8 339.87	Ed4 –Ed2.8 0.17
Clear Sky TOA SW Insolation TOA SW Up	Ed4 340.04 53.41	Ed2.8 339.87 52.50	Ed4 –Ed2.8 0.17 0.91 (1.73%)
Clear Sky TOA SW Insolation TOA SW Up TOA LW Up	Ed4 340.04 53.41 268.13	Ed2.8 339.87 52.50 265.59	Ed4 –Ed2.8 0.17 0.91 (1.73%) 2.54
Clear Sky TOA SW Insolation TOA SW Up TOA LW Up SFC SW Down	Ed4 340.04 53.41 268.13 243.72	Ed2.8 339.87 52.50 265.59 244.06	Ed4 –Ed2.8 0.17 0.91 (1.73%) 2.54 -0.33
Clear Sky TOA SW Insolation <i>TOA SW Up</i> <i>TOA LW Up</i> SFC SW Down SFC SW Up	Ed4 340.04 53.41 268.13 243.72 29.81	Ed2.8 339.87 52.50 265.59 244.06 29.74	Ed4Ed2.8 0.17 0.91 (1.73%) 2.54 -0.33 0.07
Clear Sky TOA SW Insolation <i>TOA SW Up</i> <i>TOA LW Up</i> SFC SW Down SFC SW Up SFC LW Down	Ed4     340.04     53.41     268.13     243.72     29.81     314.07	Ed2.8 339.87 52.50 265.59 244.06 29.74 316.27	Ed4Ed2.8 0.17 0.91 (1.73%) 2.54 -0.33 0.07 -2.20

### Pattern 1. SFC energy in = 2 × TOA LW out

Clear-sky	Ed2.8
TOA SW in	339.87
TOA SW up	52.50
TOA LW up	265.59
SFC SW down	244.06
SFC SW up	29.74
SFC SW in (down – up)	214.32
SFC LW down	316.27
SFC SW + LW absorbed	530.59
SFC LW up	398.40
SFC SW + LW net	132.19
G = SFC LW up – TOA LW up	132.81
2TOA LW up	531.18
Diff	-0.59

Clear-sky, Ed2.8 Surface energy absorbed SW + LW (Wm<sup>-2</sup>):

(SW down – SW up) + LW down = (244.06 – 29.74) + 316.27 = 214.32 + 316.27 = 530.59

> TOA LW out = 265.59 2 × TOA LW out = = 531.18

Diff = -0.59 Wm<sup>-2</sup>

214.32 + 316.27 = 2 × 265.59 - 0.59

#### Clear-sky, Ed2.8, time period 2001-2015, climate year:

	SW down	SW up	SW abs	LW abs	E(SFC)	OLR	20LR	Diff
CLIM 1	255.35	31.34	224.01	306.27	530.28	262.43	524.86	5.42
CLIM 2	251.86	29.86	222	307.98	529.98	262.78	525.56	4.42
CLIM 3	246.89	30.09	216.8	311.09	527.89	263.42	526.84	1.05
CLIM 4	242.26	31.63	210.63	315.14	525.77	265.01	530.02	-4.25
CLIM 5	237.33	32.21	205.12	320.27	525.39	267.18	534.36	-8.97
CLIM 6	233.41	28.67	204.74	325.70	530.44	269.05	538.1	-7.66
CLIM 7	231.63	25.72	205.91	328.35	534.26	269.75	539.5	-5.24
CLIM 8	233.65	24.42	209.23	327.05	536.28	269.12	538.24	-1.96
CLIM 9	239.24	25.80	213.44	321.82	535.26	267.58	535.16	0.1
CLIM 10	247.06	29.92	217.14	315.28	532.42	265.25	530.5	1.92
CLIM 11	253.97	33.73	220.24	309.57	529.81	263.26	526.52	3.29
CLIM 12	256.35	33.52	222.83	306.74	529.57	262.38	524.76	4.81
Average	244.08	29.74	214.34	316.27	530.61	265.60	531.20	-0.59

214.34 +316.27 = 530.61 = 2 × 265.60 - 0.59

E(SFC in, clear-sky) = (SW down – SW up) + LW in = 20LR – EEI

CERES 26th STM October 2016, Martin Wild, slide #36



TOA and SFC fluxes from independent data sources

### CERES 26th STM October 2016, Martin Wild, slide #36



solar absorbed surface + thermal down surface = 2 × thermal outgoing TOA



Energy (surface in, clear sky) = SW $\downarrow$  abs + LW $\downarrow$  abs = 2 × OLR



Surface energy budget, clear-sky, CERES EBAF Ed2.8

Solar absorbed + thermal absorbed = 2 outgoing longwave TOA  $214.34 + 316.28 = 2 \times 265.6 - 0.58 \text{ Wm}^{-2}$ SW abs + LW abs = 20LR (clear) - 0.58 Wm^{-2} => Net planetary imbalance for July 2005-June 2010: 0.58±0.43 Wm^{-2}

### Pattern 2. SFC Net = G

Clear-sky	Ed2.8
TOA SW in	339.87
TOA SW up	52.50
TOA LW up	265.59
SFC SW down	244.06
SFC SW up	29.74
SFC SW in	214.32
SFC LW in	316.27
SFC SW + LW absorbed	530.59
SFC LW up	398.40
SFC Net	132.19
G	132.81
Diff	-0.62

#### SFC Net Flux (non-radiative)

= SFC (SW in + LW in) - SFC LW up

**SFC Net** = 214.32 + 316.27 - 398.40 = **132.19** 

G = SFC LW up – TOA LW up = ULW – OLR = = 132.81

Diff (W m<sup>-2</sup>) = -0.62

### CERES 26th STM October 2016, Martin Wild, slide #36



TOA and SFC fluxes from independent data sources





G / OLR / ULW / E(SFC) = 133 / 266 / 399 / 532 = 1 / 2 / 3 / 4

### Clear-sky, Ed2.8, climate year, 2001-2015

	ULW	OLR	G	Net SFC	ULW+G	20LR	Diff
CLIM 1	388.3	262.43	125.87	141.98	514.17	524.86	10.69
CLIM 2	389.89	262.78	127.11	140.09	517	525.56	8.56
CLIM 3	393.31	263.42	129.89	134.57	523.2	526.84	3.64
CLIM 4	398.51	265.01	133.5	127.26	532.01	530.02	-1.99
CLIM 5	403.29	267.18	136.11	122.09	539.4	534.36	-5.04
CLIM 6	407.64	269.05	138.59	122.8	546.23	538.1	-8.13
CLIM 7	409.1	269.75	139.35	125.17	548.45	539.5	-8.95
CLIM 8	407.83	269.12	138.71	128.46	546.54	538.24	-8.30
CLIM 9	403.85	267.58	136.27	131.41	540.12	535.16	-4.96
CLIM 10	397.76	265.25	132.51	134.65	530.27	530.5	0.23
CLIM 11	392.27	263.26	129.01	137.54	521.28	526.52	5.24
CLIM 12	389.00	262.38	126.62	140.57	515.62	524.76	9.14
average	398.39	265.60	132.80	132.22	531.19	531.20	0.01

ULW – G = OLR (def.); **Data**: G = Net SFC (= SH+LH) ULW + G = 2OLR, Diff = 0.01 (!!!) W/m<sup>2</sup> => G = OLR/2 <=> g = G/ULW = 1/3

Greenhouse effect and normalized greenhouse factors,

years 2001 – 2015, g = G/ULW.

Integer ratios: g(all) = 6/15 = 2/5 = 0.4; g(clear) = 5/15 = 1/3

g(all-sky) and g(clear-sky), CERES EBAF Edition 2.8 (March 27, 2015), monthly mean

Theoretical lattice state at g(all-sky) = 6/15 = 2/5 = 0.4 and g(clear-sky) = 5/15 = 1/3 = 0.3333

Best fit: g(all-sky) = 0.40006 (in year 2015) g(clear-sky) = 0.33338 (in year 2011)

Increase: g(all-sky) from 0.397 to 4.000; g(clear-sky) from 0.3313 to 0.3355

2001		G = ULW - OLR	g = G/ULW		
ULW	OLR(all-sky)	G(all-sky)	g(all-sky)	OLR(clear-sky)	g(clear-sky)
386.52	236.38	150.14	0.378000	262.3	0.321381
387.61	236.47	151.14	0.375000	262.54	0.32267
391.6	236.94	154.66	0.381000	263.06	0.328243
397.01	237.91	159.1	0.390000	265.06	0.332359
402.96	240.44	162.52	0.399000	267.65	0.33579
405.53	241.76	163.77	0.406000	268.82	0.337114
407.84	243.54	164.3	0.413000	269.85	0.338343
407.41	244.1	163.31	0.416000	269.77	0.337841
403.6	241.67	161.93	0.417000	267.36	0.337562
398.07	239.21	158.86	0.399000	265.59	0.332806
392.62	237.55	155.07	0.395000	264.09	0.327365
388.71	236.21	152.5	0.392000	262.55	0.324561
Average	239.35	158.11	0.396754	265.72	0.331336

2010	OLR(all)	G	g(all)	OLR(clear)	g(clear)
388.1	237.59	150.51	0.387812	262.85	0.322726
389.76	237.47	152.29	0.390728	262.85	0.325611
393.82	238.31	155.51	0.394876	264.01	0.329618
399.41	238.16	161.25	0.40372	264.94	0.336672
404.04	240.59	163.45	0.404539	267.39	0.338209
407.37	242.28	165.09	0.405258	268.59	0.340673
408.44	243.36	165.08	0.404172	269.23	0.340833
407.45	243.86	163.59	0.401497	268.8	0.340287
403.91	242.07	161.84	0.400683	267.34	0.33812
398.44	239.44	159	0.399056	265.16	0.334505
392.67	237.56	155.11	0.395014	263.27	0.329539
387.61	236.18	151.43	0.390676	261.87	0.324398
			0.398169		0.333433
2011	OLR(all)	G	g(all)	OLR(clear)	g(clear)
386.56	235.86	150.7	0.389849	261.37	0.323857
388.4	237.52	150.88	0.388465	262.45	0.324279
392.56	237.85	154.71	0.394105	262.84	0.330446
398.11	238.17	159.94	0.401748	264.42	0.335812
403.06	239.29	163.77	0.406317	266.37	0.339131
407.06	241.97	165.09	0.405567	268.68	0.33995
408.41	243.95	164.46	0.402684	269.79	0.339414
407.51	244.05	163.46	0.401119	268.91	0.340114
403.66	242.92	160.74	0.398206	267.47	0.337388
398.28	239.02	159.26	0.399869	264.73	0.335317
391.79	236.81	154.98	0.395569	262.76	0.329335
388.35	235.99	152.36	0.392327	261.92	0.325557
			0.397985		0.33338

# Pattern 3. Clear-sky integer ratios

Costa and Shine (2012) Line-By-Line

•	ULW	= 386	6 Wm⁻²	JUNE 20	0.5 (a)		0.5	· ,	1867
•	OLR	= 259	9 Wm <sup>-;</sup>	2	(0.4 (c) (c) (c) (c) (c) (c) (c) (c) (c) (c)	M.	0.4		
•	ATM	= 194	↓ Wm <sup>-/</sup>	2	U.1 0 500	1000 1500	2000 0.1	800 1000 1200 1400	
•	G	= 127	7 Wm <sup>⊥</sup>	2	FIG. 1. Spectral dist	$B_v(T_{sfc})OLR_{clr}$ ribution of the clear-sky	STI <sub>ck</sub> (cont) ST	$\label{eq:relation} \begin{split} & \text{TI}_{ch'}(w/o \text{ cont}) \\ & \\ & \text{mponents } [W \text{ m}^{-2} \ (cm^{-1})^{-1}] \text{ using} \end{split}$	
•	STI	= 65	Wm <sup>-2</sup>		a global-mean atmosph outgoing longwave rad [STI <sub>dr</sub> (cont)]. (b) As transmitted irradiance	ere, (a) Longwave irradian liation (OLR <sub>clr</sub> ), and surfa in (a), over a smaller wav when the water vapor cont	ce emitted by surface $B_p(T_{stc})$ ce transmitted irradiance in venumber interval, but inclu inuum is excluded [STI <sub>ctr</sub> (w/c	c) assuming it to be a blackbody, the cluding the water vapor continuum des, instead of OLR <sub>elr</sub> , the surface o cont)].	
			STI	G	ATM	OLR	ULW	20LR	
	CS12	2 =	65	127	194	259	386	518	
	Patte	rn	65 /	130 /	195 /	260	/ 390 /	520	
	Ratio	OS	1 /	2 /	3 /	4 /	6 /	8	
	Di	ff	0	3	1	1	4	2 (Wm <sup>-2</sup> )	)

# E(SFC, clear) = 2OLR(clear) G(clear) = SFC Net (clear) G(clear) = OLR(clear)/2

These are **NOT** universal planetary rules.

- They cannot be deduced from the known energy in = energy out balance requirements
- They describe a unique, very specific state
- They are far from being valid, for example, on the Mars:
- The Martian ULW is 123 Wm<sup>-2</sup>, OLR = 110, G = 13 Wm<sup>-2</sup>, therefore ULW + G << 20LR ULW – OLR << 20LR – ULW G << 0LR/2.</li>



#### Read et al. (2015) QJRMS, our additions in textboxes

Energy (surface, Mars) = SW in + LW in << 20LR; 2G = 26 W/m<sup>2</sup> << OLR

# They belong to a specific geometry

 It is like the IR-opaque limit: a planet surrounded by a

> SW-transparent LW-opaque non-turbulent

"glass-shell" atmosphere.

• The surface radiation here is exactly **twice** the outgoing longwave radiation because of the construction:



# After Mashall and Plumb (2008, Fig. 2.7) **SW-transparent, LW-opaque, non-turbulent**



# All-sky

All-sky	Ed2.8
TOA SW in	339.87
TOA SW up	99.62
TOA LW up	239.60
SFC SW down	186.47
SFC SW up	24.13
SFC SW in	162.34
SFC LW down	345.15
SFC SW + LW absorbed	507.49
SFC LW up	398.27
SFC Net	109.22
G	158.67
SFC LWCRE	28.88
2TOA LW Up + SFC LWCRE	508.08
Diff	-0.59

# **Ed2.8**

#### SFC energy in:

SW in = 162.35 W/m<sup>2</sup> LW in = 345.15 W/m<sup>2</sup> SFC (SW in + LW in) = **507.5** 

#### SFC energy out:

LW up + Net = = **398.3 + 109.2** 

20LR = 2 × 239.6 = **479.2** W/m<sup>2</sup>

Diff = 507.5 - 479.2 = **28.3** W/m<sup>2</sup>

# 28 W/m<sup>2</sup> difference...

- How much was the cloud longwave radiative effect (LWCRE)?
- LWCRE = 28 W/m<sup>2</sup>.
- What we have here it is this: SFC energy in (SW + LW) = = 20LR(all) + SFC LWCRE.
- Now THAT might be meaningful :
- The surface energy budget in the all-sky has the same form as in the clear-sky case, PLUS one LW cloud radiative effect.

CLIMYEAR	SW abs	LW abs	E(SFC)	OLR(all)	LWCRE	20LR+LWCRE	Diff
CLIM 1	166.24	335.34	501.58	236.77	29.51	503.05	-1.47
CLIM 2	167.41	337.07	504.48	237.35	29.58	504.28	0.2
CLIM 3	166.63	339.65	506.28	237.72	28.92	504.36	1.92
CLIM 4	162.87	344.47	507.34	238.42	29.61	506.45	0.89
CLIM 5	157.84	349.06	506.9	240.24	28.64	509.12	-2.22
CLIM 6	155.25	353.3	508.55	242.38	28.25	513.01	-4.46
CLIM 7	156.65	355.73	512.38	243.71	28.18	515.6	-3.22
CLIM 8	160.25	355.1	515.35	243.66	28.23	515.55	-0.2
CLIM 9	163.49	350.65	514.14	242.11	28.5	512.72	1.42
CLIM 10	163.98	345.31	509.29	239.61	28.98	508.2	1.09
CLIM 11	163.53	339.77	503.3	237.15	29.86	504.16	-0.86
CLIM 12	164.09	336.33	500.42	236.28	29.75	502.31	-1.89
average	162.35	345.15	507.50	239.62	29.00	508.23	-0.73

### Pattern 4. E(SFC) = 20LR(all) + LWCRE

SFC (SW in + LW in) = 20LR(all) + LWCRE (-0.73 W/m<sup>2</sup>)



 $E(SFC, all) = LW in + SW in = 2OLR(all) + LWCRE + IMB - 0.05 (!!!) W/m^2$ 344 + 163 = 2 x 240 + 26.6 + 0.45 - 0.05

### All-sky

All-sky	Ed2.8
TOA SW in	339.87
TOA SW up	99.62
TOA LW up	239.60
SFC SW down	186.47
SFC SW up	24.13
SFC SW in	162.34
SFC LW down	345.15
SFC SW + LW absorbed	507.49
SFC LW up	398.27
SFC Net	109.22
G	158.67
SFC LWCRE	28.88
2TOA LW Up + SFC LWCRE	508.08
Diff	-0.59

### Ed2.8

=

# SFC energy in

(SW + LW ) = = 162.34 + 345.15 = **507.49** 

2 x TOA LW out + SFC LWCRE = 2 x 239.6 + 28.88 = 508.08

Diff = -0.59 W m<sup>-2</sup>

# Ed4.0

All-sky	Ed4.0
TOA SW in	340.04
TOA SW up	99.23
TOA LW up	240.14
SFC SW down	187.04
SFC SW up	23.37
SFC SW in	163.67
SFC LW down	344.97
SFC SW + LW absorbed	508.64
SFC LW up	398.34
SFC Net	110.30
G	158.20
SFC LWCRE	30.90
2TOA LW Up + SFC LWCRE	511.18
Diff	-2.54

### All-sky, Ed4.0

Energy absorbed SFC (W m<sup>-2</sup>):

SW in + LW in = 163.67 + 344.97 = **508.64** 

2 x OLR + SFC LWCRE = 2 x 240.14 + 30.90 = **511.18** 

Diff = -2.54 W m<sup>-2</sup>

# Ed4.0

Clear-sky	Ed4.0
TOA SW in	340.04
TOA SW up	53.41
TOA LW up	268.13
SFC SW down	243.72
SFC SW up	29.81
SFC SW in	213.91
SFC LW down	314.07
SFC SW + LW Absorbed	527.98
SFC LW Up	397.59
SFC Net	130.39
G	129.46
Diff	0.93

Clear-sky, Ed4.0 SFC Net = G

SFC Net = SW in + LW in – LW up = 528.0 – 397.6 = **130.4** W m<sup>-2</sup>

G = ULW – OLR = 397.6 – 268.1 = **129.5** W m<sup>-2</sup>

Diff = **0.9** W m<sup>-2</sup>.

# Ed4.0 Clear-sky: SFC Net = G

	LILW clear	OLP alaar	Galaar	SEC Not	Diff
	ULVV Clear	ULK Clear	G clear	SFC Net	DITT
CLIM 1	388.08	265.08	123	139.01	16.01
CLIM 2	389.63	265.27	124.36	137.49	13.13
CLIM 3	393.06	266.02	127.04	132.44	5.4
CLIM 4	397.64	267.04	130.6	126.25	-4.35
CLIM 5	402.18	268.99	133.19	121.41	-11.78
CLIM 6	405.9	271	134.9	122.11	-12.79
CLIM 7	407.29	272	135.29	124.59	-10.7
CLIM 8	406.4	271.71	134.69	127.49	-7.2
CLIM 9	402.83	270.36	132.47	129.98	-2.49
CLIM 10	397.57	268.14	129.43	132.33	2.9
CLIM 11	392	265.95	126.05	134.61	8.56
CLIM 12	388.69	264.91	123.78	137.19	13.41
Average	397.606	268.039	129.57	130.41	0.84

G= ULW - OLR= 397.6 - 268.0= 129.6 W m<sup>-2</sup>Diff== 0.8 W m<sup>-2</sup>

### Pattern 5. All-sky integer ratios





### $F = N \times UNIT$ ; UNIT = OLR(all-sky)/9



# Is it possible to satisfy all the patterns with one data set?

Let's try.

# Model data set: EdMZ All-sky pattern positions

All-sky	Ed4.0	Ed2.8	EdMZ	Ed4.0 – EdMZ	Ed2.8 – EdMZ
TOA SW In	340.04	339.87	340.04	0.0	-0.27
TOA LW Up	240.14	239.60	240.14	0.0	-0.54
SFC SW In	163.67	162.34	160.09	3.58	2.25
SFC LW Down	344.97	345.15	346.87	-1.90	-1.72
SFC (SW in + LW in)	508.64	507.49	506.96	1.68	0.53
SFC LW Up	398.34	398.27	400.23	-1.89	-1.96
SFC Net	110.30	109.22	106.73	3.57	2.49
G	158.20	158.67	160.09	-1.89	-1.42
2TOA LW Up+LWCRE	511.18	508.08	506.96	4.22	1.12

## EdMZ all-sky integer ratios F = N × UNIT UNIT = OLR(all-sky)/9

All-sky Flux	EdMZ	N
TOA SW In	340.04	
TOA SW Up	99.60	
TOA LW Up	240.14	9
SFC SW In	160.09	6
SFC LW Down	346.87	13
SFC (SW in + LW in)	506.96	19
SFC LW Up	400.23	15
SFC Net	106.73	4
G	160.09	6
SFC LWCRE	26.68	1
2 × TOA LW Up + LWCRE	506.96	19

# Model data set: EdMZ Clear-sky pattern positions

Clear-sky	Ed4.0	Ed2.8	EdMZ	Ed4.0 – EdMZ	Ed2.8 – EdMZ
TOA SW in	340.04	339.87	340.04	0.0	-0.17
TOA LW up	268.13	265.59	266.82	1.31	-1.23
SFC SW in	213.91	214.32	213.47	0.44	0.85
SFC SW + LW in	527.98	530.59	533.65	-5.67	-3.06
SFC LW up	397.59	398.40	400.23	-2.64	-1.83
SFC Net	130.39	132.19	133.42	-3.03	-1.23
G	129.46	132.81	133.42	-3.96	-0.61

### EdMZ clear-sky integer ratios F = N × UNIT UNIT = OLR(clear-sky)/4

Clear-sky Flux	EdMZ	N
TOA SW In	340.04	
TOA LW Up	266.82	4
ATM emitted Up	200.11	3
STI	66.7	1
SFC (SW + LW) In	533.65	8
SFC LW Up	400.23	6
SFC Net	133.42	2
G	133.42	2

# Clear-sky fluxes in all-sky units F = N × UNIT(all-sky)

Clear-sky Flux	EdMZ	N
TOA SW In	340.04	
TOA LW Up	266.82	10
SFC SW In	213.47	8
SFC LW Down	320.18	12
SFC (SW + LW) In	533.65	20
SFC LW Up	400.23	15
SFC Net	133.42	5
G	133.42	5
TOA LWCRE (UNIT)	26.68	1

# The patterns as integers



CLEAR SKY basics: g = (15 – 10) /15 = **5/15** = 1/3









Solar Absorbed Surface (SAS) = 3 goes into G = ULW - OLR = 3



**Step 2** Allow **ONE** atmospheric SW-absorption: SAA = 1, SAS = 2

Solar Absorbed Atmosphere (SAA) = 1, Solar Absorbed Surface (SAS) = 2









SAS (6) = G (6)

Step 5 Allow ONE partial LW-transparency ...



Step 6 ... fade the window with ONE up and down LW cloud effect



Step 7 ... and close the balance with turbulence. Atmosphere: E(SFC) + 2 UNITS = 21 = emitted up (8) + down (13)



Surface: E(SFC) = 2 OLR + 1 UNIT = 19

The pattern. Basic energy flow routes and integer rates.







# EdMZ with ISR = 340.0 and OLR = 240.1 Wm<sup>-2</sup>



# Specific geometries

Moon (zero)			
ULW – G = OLR	G = 0	g = 0	
ULW + G = OLR	ULW = OLR	t = 1	
ז	Mars (intermediate, free)		
ULW – G = OLR	G = 0.118 OLR	g = 0.106	
ULW + G = 1.236 OLR	ULW = 1.118 OLR	t = 0.732	
Ea	arth (clear-sky) (level one)		
ULW – G = OLR	G = (1/2) OLR	g = 1/3	
ULW + G = 20LR	ULW = (3/2) OLR	t = 1/6	
Earth (all-sky) (level two)			
ULW – G = OLR	G = (2/3) OLR	g = 2/5	
ULW + G = (7/3) OLR	ULW = (5/3) OLR	t = 1/15	
Shield (level three)			
ULW – G = OLR	G = OLR	g = 1/2	
ULW + G = 30LR	ULW = 20LR	t = 0	

t = STI/ULW, atmospheric LW transmittance

**Sw The abed of Earth Reviews of Geophysics** Graeme L. Stephens<sup>1,2,3</sup>, Denis O'Brien<sup>4</sup>, Peter J. Webster<sup>5</sup>, Peter Pilewski<sup>6,7</sup>, Seiji Kato<sup>8</sup>, and Jui-lin Li<sup>1</sup> "surprising" hemispheric symmetry and "remarkable" interannual stability of the system albedo. *Why?* They refer to **Gaia**. I think the **Blue Marble** (Glass Shell) is better:  $\alpha_0 = 1 - \sin 45^\circ$   $= 1 - \sqrt{2/2}$ = 0.292893

CERES - EBAF Data Time Range: January to December - CLIMATE YEAR Time Resolution: CLIM Valid Range: 0 - 800

Incomi	.ng	Solar Flux	TOA Shortwave Flux	Albedo
(W m-2	2)		All-Sky (W m-2)	
	121			
CLIM	1	350.69	106.26	0.30300265
CLIM	2	348.15	102.27	0.29375269
CLIM	3	343.98	99.31	0.28870865
CLIM	4	337.54	97.44	0.2886769
CLIM	5	332.3	97.45	0.2932591
CLIM	6	329.24	96.39	0.29276516
CLIM	7	328.89	93.93	0.28559701
CLIM	8	331.43	92.62	0.27945569
CLIM	9	336.78	94.79	0.28145971
CLIM	10	342.04	100.24	0.29306514
CLIM	11	347.05	106.18	0.30595015
CLIM	12	350.34	108.29	0.30909973
AVERA	GE	339.8692	99.5975	0.29289938

# Climate sudoku







### $1 = 26.6 \text{ W/m}^2$ , $\Delta \max(\text{Wild-EdMZ}) = -3.8 \text{ W/m}^2 (\text{DLR})$

# Summary

- There are robust patterns in the annual global means.
- EdMZ is an idealized data set representing the pattern, belonging to a time-independent geometry.
- The integer ratios follow from the closed-shell geometry.
- The corresponding physical state has some reasonable theoretical basis.
- The largest bias between Ed4.0 and EdMZ is -6.1 Wm<sup>-2</sup> (2%) in DLR(clear) and 3.6 Wm<sup>-2</sup> (2.2%) in SFC SW(all).
- Size and time-scale of fluctuations around or systematic deviations from the pattern positions are not yet known.





# 30 Wm<sup>-2</sup> CMIP5 model projected increase in DLR

(c) Changes in surface downward longwave radiation (Wm<sup>-2</sup>)



Stephens et al. (2012) Nat Geosci Suppl Fig S2 (c)

# 30 Wm<sup>-2</sup> increase in DLR would falsify EdMZ



