


## 1-minute summary

- Backprojection (BP) is a powerful seismic-array processing tool that provides observational constraints on the evolution of the seismic radiation during large earthquakes. To better interpret the BP images in terms of earthquake dynamics, a careful relation between the BP images and the kinematics of the source is needed.
- We explore the relation between BP images and seismic radiation using synthetic waveforms from both kinematic and dynamic source models embedded in a pure elastic wholespace.
- Preliminary results show the BP image is mostly correlated to the distribution of peak slip/moment rate, for both kinematic and dynamic sources. However, it is also highly related to other factors thus should be carefully interpreted with coseismic observations.

## Methodology

### 1. Source models

- Kinematic source models are generated by FFSP (source generator developed by Liu *et al.* [2006] and Crempien and Archuleta [2014])
- Dynamic source model is from Dr. Huihui Weng for the 2015 Gorkha Mw 7.8 earthquake based on slip weakening with parameters constrained by near-field ground motion observations.



### 4. 2D Image correlation (quantified comparisons)

### 3. BP in frequency domain

$$B(f) = A(f)X(f)$$

$$X(f) = A^H(f)B(f)/N$$


$B(f)$ : Velocity data in frequency domain  
 $A(f)$ : Propagation matrix (travel time shifting)  
 $X(f)$ : Distribution of BP power  
 $H$ : Conjugate transpose  $N$ : Number of stations

### 2. Synthetic waveforms:


Waveforms are derived from the slip histories of the source models in the homogeneous, elastic wholespace.

## Results

### 1. BP image & Kinematic sources




↑ 2-4 Hz integrated BP image (black contours) is most correlated to average moment rate, spatial gradient (strain/stress rate?) and rupture velocity variation.



↑ Correlations with source parameters for all 18 generated kinematic sources in the integrated frequency band of 2-4Hz.

- Correlations (crosses) with source parameters for all sources in each narrow frequency band. Correlation coefficients vary for different sources and there is no systematic frequency dependence on the correlation coefficients of each source parameters.
- Average correlation coefficients (red lines) seem to be relatively stable, implying the general correlation between BP image and all these parameters.

### 2. BP image & Dynamic source model of 2015 Nepal Gorkha Mw 7.8 earthquake



↑ Correlation between BP image and dynamic model within 2-4Hz

Peak moment rate > Moment > Rupture velocity (?) > Stress drop ≈ Initial stress > Rise time

➤ Correlations with source parameters from the dynamic source in each narrow frequency band. Systematic frequency dependence on the correlation coefficients is unclear for this dynamic model, the valley around 2.8Hz is mostly due to swimming artifacts of conventional BP.

## Conclusion and Discussion

- Both kinematic and dynamic source models indicate correlations between BP image and distributions of slip, slip rate, slip gradient (strain), slip rate gradient (strain rate) and rupture velocity heterogeneities.
- No single parameter dominates the characteristics of BP images, but rather a combination of them (moment, moment rate).
- Our results show no systematic frequency-dependent correlation patterns, which implies that the differences reported in BP images with variable seismic frequencies probably stem from other factors such as fault geometry, roughness and/or frictional properties etc.

### Future work:

- Apply to more diverse kinematic source models for better parametrization (i.e. varying asperity size and amplitude of heterogeneity)
- Use advanced BP methods to gain better BP image with less artifacts

### Acknowledgements:

We sincerely thank Pengcheng Liu, Jorge Crempien and Ralph Archuleta for the FFSP code and Huihui Weng for the Nepal dynamic rupture model. More detailed discussion about the dynamic rupture model can be found in poster (S51A-0580).