



# Seismic Evidence for Partial Melt Below Tectonic Plates



Eric Debayle<sup>(1)</sup>, Thomas Bodin<sup>(1)</sup>, Stéphanie Durand<sup>(1)</sup>, Yanick Ricard<sup>(2)</sup>  
(1) Univ Lyon, Univ Lyon 1, ENSL, CNRS, LGL-TPE, Villeurbanne, France - (2) Univ Lyon, ENSL, Univ Lyon 1, CNRS, LGL-TPE, Lyon, France.



PRESENTED AT:  
**AGU FALL MEETING**  
New Orleans, LA & Online Everywhere  
13-17 December 2023  
Poster Gallery  
WILEY

**ABSTRACT :**  
The paper related to this research is available from [nature website](#).

Observations of shear attenuation provide new constraints on the origin of the seismic low-velocity zone (LVZ) in the upper mantle. We report the presence of partial melt in the LVZ, from the joint interpretation of 3D shear attenuation and velocity models. We observed that partial melting up to 10–30 kilometres deep feeds the asthenosphere beneath mid-ocean ridges, major hotspots and back-arc regions. A small amount of melt (less than 0.3%) remains trapped within the oceanic LVZ. Melt is mostly absent under continental regions. The amount of melt increases with plate velocity, particularly when the plate velocities change from 3 cm/year to 5 cm/year. This finding is consistent with previous observations of mantle crystal alignment underneath tectonic plates. Our observations suggest that by reducing viscosity, melt facilitates plate motion and large-scale crystal alignment in the asthenosphere.

## FIFTY YEARS OF DEBATE ON THE ORIGIN OF THE ASTHENOSPHERE :

**Arguments supporting the presence of melt in the LVZ :**

- Simple idea that explains the Vs reduction and the low viscosity required in the LVZ (Anderson and Sammis, 1970)
- Production of MORB by partial melting is the largest volcanic activity on Earth
- Difficulty to explain the sharp Vs contrasts with purely thermal model (Kawakatsu et al. 2009)
- Surface tension may resist complete draining of small melt fraction in the LVZ (Holtzman, 2016)

**Arguments against the presence of melt :**

- The Vs reduction in the LVZ can be explained by solid state mechanisms without the presence of melt or fluids (Faul and Jackson, 2005)
- Small melt fractions (<10<sup>-3</sup>% volume) may move upward the lithospheric plate above and will solidify (Mc Kenzie, 1989)
- The sharp velocity reduction associated with the LAB and MLD may be explained by Elastically Accommodated Grain boundary Sliding (EAGBS), a subolidus deformation mechanisms (Karato, 2012, Karato et al., 2015)

It is difficult to constrain the origin of the LVZ with seismic velocity alone. Adding quality factor provides new independent constraints!

## DATASETS AND METHODOLOGY

**Dataset:**  
We use 2 consistent Vs and Qs 3D models, D2020s and QsADRI1 are :

Based on the same dataset, Debayle and Ricard, (2012) fundamental and higher modes global dataset (372,629 Rayleigh waveforms)

Inverted using the same approach with the same horizontal and vertical smoothing

**Methodology :**  
We plot Vs as a function of ln(Qs) for each geographical point and compare our observations with predictions from :


- a temperature dependent model (Takei, 2017) - dark blue curve
- experimental results from Jackson et al., (2002) - light blue curve

assuming a pyroclitic mantle.

The blue curves do not explain all data. Adding melt reduces velocities and shift the blue curves to the left. The color scale indicates the amount of melt (%) required to explain data at each geographical point. We prefer Takei's model (dark blue curve) as it minimizes the required amount of melt.

The effect of melt on Vs is predicted using experimental results from Chantel et al., 2016 (red curve)


We neglect the effect of melt on Qs for long period surface waves for the following reasons : The Melt experiment results (Yang et al., 2007) suggest that for long period surface waves (>20 s) melt has a very small effect on Qs. This is also expected if the attenuation mechanism is melt squirt (Hammond and Humphreys, 2006).



The figure above (left) shows range of acceptable Qs average after Yang et al., (2007) for a 2 Ma seafloor (red boxes), a 6 Ma sea floor (orange boxes) and the average oceanic values (about 80 Ma, blue boxes).

## MELT CONTENT AND EFFECT ON VISCOSITY


Melt content maps are displayed at different depths in the upper mantle.



The upper colour scale shows the amount of melt (%) required to explain our observations using the model of Takei, (2017). The lower colour scale indicates the difference in percent between theory and observations, in regions (mostly contours) where Vs is too high and cannot be reconciled with model predictions assuming a pyroclitic mantle. In regions (mostly contours) where Vs is too low and cannot be reconciled with model predictions assuming a pyroclitic mantle, the difference is indicated with black triangles. The blue and green stars indicate the location of the North (Lin et al., 2016) and Melt (Yang et al., 2007) experiments, respectively.


We map partial melting down to 150–200 kilometres beneath mid-ocean ridges, major hotspots and back-arc regions. In these regions melt feeds the asthenosphere. A small part of this melt (less than 0.3%) remains trapped within the oceanic LVZ. Melt is mostly absent under continental regions, where compositional heterogeneities and depletion of incompatible elements contribute to the high seismic velocities.

**Effect of melt on viscosity :**  
The figure below by Holtzman (2016) shows that the variation in viscosity as a function of melt content occurs in two steps. First, the onset of melting has a strong effect on viscosity when a connected network of melt tubules is established. The viscosity decreases by one or two orders of magnitude before the melt fraction  $\phi$  reaches 0.1%. Second, for larger melt content, the viscosity decreases further, with  $\ln(\eta/\eta_0) \sim -2\phi$ , this effect is minor for the low melt content that we observe. The viscosity should be one or two orders of magnitude lower under oceanic plates, where  $\phi < 0.3\%$ , than under continents.



## EFFECT OF COMPOSITION AND RADIAL ANISOTROPY

**Composition:** We assumed the mantle is pyroclitic. Can we explain our observations with a change of composition?




We test 3 extreme compositions: an upper mantle entirely made of harzburgite (blue box and curves), of silica-excess pyroxenite (orange box and curves) or of silica deficient pyroxenite (red boxes and curves).

Neither of these extreme compositions can remove the melt layer in the LVZ.


**Radial anisotropy:** can we explain our observations without melt, only by invoking radial anisotropy (displayed with  $\xi = (V_{sh}/V_{sv})^2$ )

The figure below shows the global average of the radial anisotropy (blue curve) needed to explain our observations without melt. The  $\xi$  values are unrealistic and much larger than the global average in 3D models (S3M120 (Vinnikov et al., 2013, green curve), S3M21 (Kawakatsu et al., 2008, brown curve) or PREM (Dziewonski & Anderson, 1981, red curve).



We show in the Method and Supplementary information of our paper that the low Vs value cannot be reconciled with Qs by adding radial anisotropy, elastically accommodated grain-boundary sliding, or the effect of composition or water.

## THE AMOUNT OF MELT INCREASES WITH PLATE VELOCITY



Melt fraction at different depths (left panel) and anisotropy along asthenotic plate motion (right panel, continuous lines) as a function of plate velocity expressed in a nonet reference frame for different depths in the upper mantle. The amount of melt increases substantially in the asthenosphere (100–200 km) for plate velocities between 3 and 5 cm/yr (left panel). This links with previous observations by Debayle and Ricard (2013) (right panel) of an increase in the proportion of anisotropy aligned with APM for plate motions between 3 and 5 cm/yr.

Plate crystal alignment beneath plates moving faster than 4 cm/yr (right panel) is associated with a greater amount of melt (left panel).

This requires either that melt facilitates deformation or that deformation favours melt retention or both.

Our results suggest that by reducing viscosity, melt facilitates plate motion and large-scale crystal alignment in the asthenosphere.

## DESIGN YOUR IPOSTER

Background:  Title Text Color:  Textbox Title Color:

Default

Fill Color:  Title Text Color:  Side Textbox Title Background Color:

Subtitle Text Font:  Textbox Text Font:

Gradient Color:  Default  Default

Subtitle Text Color:  Side Textbox Text Color:

OPEN Arrow Color:  Side Textbox Background Color:

Light Gray  Textbox Title Font:  Middle Textbox Text Color:

Default  Middle Textbox Background Color:

Notice!  
Your iPoster has now been unpublished and will not be displayed on the iPoster Gallery.  
You need to publish it again if you want to be displayed.

You must be registered to publish your poster.  
Please visit [service@agu.org](#)

Because of maintenance we have just saved your content and will within a few minutes logout all users and restart our server. We will be back in a moment.  
Sorry for the inconvenience!  
Because of maintenance we will within a few minutes restart our server. We will be back in a moment.  
Sorry for the inconvenience!

LINK:  
AUTHOR INFORMATION

Eric Debayle<sup>1</sup>, Thomas Bodin<sup>1</sup>, Stéphanie Durand<sup>1</sup>, Yanick Ricard<sup>2</sup>  
1Univ Lyon, Univ Lyon 1, ENSL, CNRS, LGL-TPE, Villeurbanne, France  
2Univ Lyon, ENSL, Univ Lyon 1, CNRS, LGL-TPE, Lyon, France.  
e-mail: [Eric.Debayle@ens-lyon.fr](mailto:Eric.Debayle@ens-lyon.fr)

## ABSTRACT

The seismic low-velocity zone (LVZ) of the upper mantle is generally associated with a low-viscosity asthenosphere that has a key role in decoupling tectonic plates from the mantle. However, the origin of the LVZ remains unclear. Some studies attribute its low seismic velocities to a small amount of partial melt of minerals in the mantle, whereas others attribute them to solid-state mechanisms near the solidus or the effect of its volatile contents. Observations of shear attenuation provide additional constraints on the origin of the LVZ. On the basis of the interpretation of global three-dimensional shear attenuation and velocity models, here we report partial melt occurring within the LVZ. We observe that partial melting down to 150–200 kilometres beneath mid-ocean ridges, major hotspots and back-arc regions feeds the asthenosphere. A small part of this melt (less than 0.30 per cent) remains trapped within the oceanic LVZ. Melt is mostly absent under continental regions. The amount of melt increases with plate velocity, increasing substantially for plate velocities of between 3 centimetres per year and 5 centimetres per year. This finding is consistent with previous observations of mantle crystal alignment underneath tectonic plates. Our observations suggest that by reducing viscosity, melt facilitates plate motion and large-scale crystal alignment in the asthenosphere.

## REFERENCES

- Anderson D.L. & C. Sammis(1970). Partial melting in the upper mantle, *Physics of the Earth and Planetary Interiors*, Volume 3, Pages 41–50.
- Debayle, E. & Ricard (2012). A global shear velocity model of the upper mantle from fundamental and higher Rayleigh mode measurements. *J. Geophys. Res. Solid Earth* 117, B10308.
- Dziewonski, A., and D. Anderson (1981). Preliminary reference Earth model, *Phys. Earth Planet. Inter.*, 25, 287–356.
- Faul, U. H. & Jackson, I. (2005). The seismological signature of temperature and grain size variations in the upper mantle. *Earth Planet. Sci. Lett.* 234, 119–134.
- French, S., Lekic, V. & Romanowicz, B. (2013). Waveform tomography reveals channelled flow at the base of the oceanic asthenosphere. *Science* 342, 227–230.
- Holtzman, B. K. (2016). Questions on the existence, persistence, and mechanical effects of a very small melt fraction in the asthenosphere. *Geochim. Geophys. Geostyst.* 17, 470–484.
- Karato, S. (2012). On the origin of the asthenosphere. *Earth Planet. Sci. Lett.* 321–322, 95–103.
- Karato, S. I., Okubo, T. & Park, (2015). J. Mechanisms and geologic significance of the mid-lithosphere discontinuity in the continents. *Nat. Geosci.* 8, 509–514.
- Kawakatsu, H. et al. (2009). Seismic evidence for sharp lithosphere-asthenosphere boundaries of oceanic plates. *Science* 324, 499–502.
- Kustowski, B., Ekström, G. & Dziewonski, A. M. Anisotropic shear-wave velocity structure of the Earth's mantle: a global model. *J. Geophys. Res.* Solid Earth 113, B06306 (2008).
- Lin, P. Y. P. et al. High-resolution seismic constraints on flow dynamics in the oceanic asthenosphere. *Nature* 535, 538–541 (2016).
- McKenzie, D. (1989). Some remarks on the movement of small melt fractions in the mantle. *Earth & Planetary Science Letters*, 95, 53–72
- Takei, Y. (2017). Effects of partial melting on seismic velocity and attenuation: a new insight from experiments. *Annu. Rev. Earth Planet. Sci.* 45, 447–470.
- Yang, Y., Forsyth, D. W. & Weeraratne, D. S. Seismic attenuation near the East Pacific Rise and the origin of the low-velocity zone. *Earth Planet. Sci. Lett.* 258, 260–268 (2007).

## CHAT SETTINGS

Choose Date & Time: Your Timezone:  Time:  Date:

Edit Chatroom Message: Hello! My chat will be starting at the time listed below. If the chat isn't open, it means I'm not here yet - so please contact me using the Contact Author button at the bottom of my iPoster.

Please note: After setting up or changing your session time, it may take up to an hour for the change to be displayed on the Gallery screen.

## YOUR CHAT SESSION SCHEDULE

DateStart	End	Time zone	Actions	<input type="button" value="Add new session"/>
-----------	-----	-----------	---------	--

## CHAT INFORMATION

## IN PERSON POSTER HALL SESSION SETTINGS

Choose Date & Time: Your Timezone:  Time:  Date:

Edit Session Message:

Session - Board Number:

Session - Board Number:

Please note: After setting up or changing your session time, it may take up to an hour for the change to be displayed on the Gallery screen.

## YOUR IN PERSON POSTER HALL SESSION SCHEDULE

Date	Start	End	Time zone	Actions	<input type="button" value="Add new session"/>
------	-------	-----	-----------	---------	--

## SHARE POSTER

SHARE LINK:

## JOIN CHAT

In order to join Chat, please enter your name and email address

Enter your name here:

Enter your email address here:

## AT MY POSTER

## REVIEWER SURVEY